

Rev04
Update: Oct,19,2022

DATASHEET

MonoRab™ HA tag Antibody(109B2), mAb, Rabbit

Cat. No.: A01963

Overview

Specificity	MonoRab™ HA tag Antibody recognizes HA tags placed at N-terminal, C-terminal, and internal regions of fusion proteins.
Host Species	Rabbit
Immunogen	a synthetic peptide containing the influenza hemagglutinin epitope (YPYDVPDYA). conjugated to KLH
Conjugate	Unconjugated

Applications

Working concentrations for specific applications should be determined by the investigators. The appropriate concentration may be affected by secondary antibody affinity, antigen concentration, the sensitivity of the method of detection, temperature, the length of the incubations, and other factors. The suitability of this antibody for applications other than those listed below has not been determined. The following concentration ranges are recommended starting points for this product.


Application	Recommended Usage
ELISA	0.05-0.2 µg/ml
Western Blot	0.1-0.2 µg/ml
Immunocytochemistry/Immunofluorescence (ICC/IF)	0.5-2 µg/ml

Properties

Form	Lyophilized
Storage Buffer	lyophilized with PBS, pH 7.4, containing 0.02% sodium azide.
Reconstitution	Reconstitute the lyophilized powder with deionized water (or equivalent) to an final concentration of 0.5 mg/mL.


Storage Instructions	The lyophilized product remains stable up to 1 year at -20 °C from date of receipt. Upon reconstitution, it can be stored for 2-3 weeks at 2-8 °C or for up to 12 months at -20 °C or below. Avoid freeze/thaw cycles.
Purification	Protein A affinity column
Isotype	Rabbit IgG
Clonality	Monoclonal
Clone ID	109B2
Note	GenScript can customize this product per customer's request including product size, buffer components, etc.

Examples



Immunofluorescence staining of MonoRab™ HA tag Antibody (109B2), mAb, Rabbit (A01963-40) with HA-tagged red fluorescent protein expression in CHO cells.

The cells were fixed with 4% Poly-Formaldehyde for 5min, permeabilized with 0.1% TritonX-100 for 5 minutes, and blocked in 3% BSA for 30min at room temperature. The cells were then stained with 1/1000 Rabbit Anti-HA-tag mAb at room temperature for 2h, followed by a further incubation at 37 °C for 1h with Mouse Anti-Rabbit IgG Antibody [iFluor488], mAb at 5 µg/ml.



Western blot analysis of MonoRab™ HA tag Antibody (109B2), mAb, Rabbit with HA-tag fusion proteins. Loading:

A: C-terminal HA-tagged fusion protein (25 ng,10 ng,5 ng)

B: N-terminal HA-tagged fusion protein (25 ng,10 ng,5 ng)

C: M-terminal HA-tagged fusion protein (25 ng,10 ng,5 ng)

Primary Antibody:

MonoRab™ HA tag Antibody(109B2), mAb, Rabbit (A01963-40) 0.2 µg/ml

Secondary Antibody:

Goat anti-rabbit IgG (H&L) [IRDye⁸⁰⁰] (Licor,926-32211) 0.125 µg/ml

Background

Target Background : The rabbit immune system generates antibody diversity and optimizes affinity. GenScript utilizes MonoRab™ technology to generate high affinity and specific monoclonal rabbit antibodies. The HA tag is composed of several amino acids containing YPYDVPDYA and does not appear to interfere with the bioactivity or the biodistribution of recombinant proteins. GenScript MonoRab™ HA tag Antibody (109B2), mAb, Rabbit is specific to HA tags placed at the N-terminal, C-terminal, and internal regions of fusion proteins. This antibody can greatly improve the effectiveness of several different kinds of immunoassays, helping researchers identify and detect HA-tagged fusion proteins in bacteria, insect cells, and mammalian cells.

Synonyms : Rabbit monoclonal to HA tag

For laboratory research use only. Direct human use, including taking orally and injection and clinical use are forbidden.