

Rev04
Update: Oct,19,2022

DATASHEET

Anti-Bevacizumab Antibody (6C3), mAb, Mouse

Cat. No.: A01976

Overview

| | |
|---------------------|--|
| Specificity | The product is specific for Bevacizumab. The antibody is recommended as a detection antibody in a pharmacokinetic (PK) bridging assay with capture antibody GenScript, A01975-40, Anti-Bevacizumab Antibody (4H1), mAb, Mouse. |
| Host Species | Mouse |
| Immunogen | Bevacizumab |
| Conjugate | Unconjugated |

Applications

Working concentrations for specific applications should be determined by the investigators. The appropriate concentrations may be affected by secondary antibody affinity, antigen concentration, the sensitivity of the method of detection, temperature, the length of the incubations, and other factors. The suitability of this antibody for applications other than those listed below has not been determined. The following concentration ranges are recommended starting points for this product.

| Application | Recommended Usage |
|--------------------|--------------------------|
| ELISA | 0.01-0.1 µg/ml |
| Competitive ELISA | 0.1-10 µg/ml |

Properties

| | |
|-----------------------------|--|
| Form | Lyophilized |
| Storage Buffer | lyophilized with PBS, pH 7.4, containing 0.02% sodium azide. |
| Reconstitution | Reconstitute the lyophilized powder with deionized water (or equivalent) to an final concentration of 0.5 mg/mL. |
| Storage Instructions | The lyophilized product remains stable up to 1 year at -20 °C from date of receipt. Upon reconstitution, it can be stored for 2-3 weeks at 2-8 °C or for up to 12 months at -20 °C or below. Avoid freeze/thaw cycles. |
| Purification | Protein A affinity column |

GenScript USA, Inc.


860 Centennial Ave. Piscataway, NJ 08854

Tel: 1-732-885-9188

| | |
|------------------|---|
| Isotype | Mouse IgG1 |
| Clonality | Monoclonal |
| Clone ID | 6C3 |
| Note | GenScript can customize this product per customer's request including product size, buffer components, etc. |

Examples

Anti-Bevacizumab Antibody (6C3) blocks Bevacizumab with VEGF165



Anti-Bevacizumab Antibody (6C3) (GenScript, A01976-40) blocks Bevacizumab binding with VEGF165 protein (GenScript, Z03073-50).


Coating antigen: VEGF165, 1 µg/ml.

Bevacizumab final concentration: 0.1 µg/ml.

Anti-Bevacizumab Antibody (6C3) (GenScript, A01976-40) dilutions start from 10 µg/ml.

IC₅₀= 0.157 µg/ml.


Anti-Bevacizumab Antibody (6C3) binds with Bevacizumab



ELISA binding of Anti-Bevacizumab Antibody (6C3) (GenScript, A01976-40) with Bevacizumab. While the antibody does not recognize the human IgG (data not shown). Coating antigen: Bevacizumab, 1 µg/ml.

Anti-Bevacizumab Antibody (6C3) (GenScript, A01976-40) dilutions start from 1,000 ng/ml.

EC₅₀= 8.735 ng/ml.



Standard curve of Bevacizumab Sandwich ELISA. The Bevacizumab Sandwich ELISA assay is developed by using Anti-Bevacizumab Antibody (4H1), mAb, Mouse (GenScript, A01975-40) and Anti-Bevacizumab Antibody (6C3), mAb, Mouse (GenScript, A01976-40) as the capture and detection antibodies, respectively

In this ELISA assay, Anti-Bevacizumab Antibody (6C3), mAb, Mouse (GenScript, A01976-40) was labeled with Biotin. GenScript can provide customized conjugation services for this product per customer's request.

The sensitivity is less than 78 pg/ml.

Background

Target Background : Bevacizumab (Avastin) is a humanized monoclonal antibody that is approved by the U.S. Food and Drug Administration for the treatment of a number of types of cancers and a specific eye disease. Bevacizumab binds a continuous epitope in VEGF, preventing interaction of VEGF with VEGFR on the surface of endothelial cells. It can be used to inhibit solid tumour growth and metastasis by weakening tumour angiogenesis. Anti-Bevacizumab Antibody (6C3), mAb, Mouse is produced from a hybridoma resulting from the fusion of partner and B-lymphocytes obtained from a mouse immunized with Bevacizumab.

Synonyms : Mouse monoclonal to Avastin

For laboratory research use only. Direct human use, including taking orally and injection and clinical use are forbidden.