

GM-CSF (Sargramostim), Human (*P. pastoris*-expressed)

Cat. No.: Z02693-1

Size: 1.0 mg

Synonyms: Granulocyte Macrophage Colony Stimulating Factor, CSF-2, MGI-1GM, GM-CSF, Pluripoietin-alpha, Molgramostin, Sargramostim, MGC131935, MGC138897

Description:

Human Granulocyte Macrophage Colony Stimulating Factor (hGM-CSF), a hematopoietic growth factor, is mainly involved in granulopoiesis and monocytopoiesis. It is produced by T-cells and macrophages in response to antigens, and by endothelial cells and fibroblasts following induction of various cytokines^[1]. A monomeric protein of 127 amino acids with six glycosylation sites and two intra disulfide bonds^[2,3], glycosylated and non-glycosylated hGM-CSFs show similar biological activities^[4]. Other than its connection to the growth and development of granulocytes and macrophages, it is also indispensable for the proliferation of erythroid and megakaryocytic cells^[5].

Recombinant Human Granulocyte Macrophage Colony Stimulating Factor (rhGM-CSF) produced in *P. pastoris* is a glycosylated polypeptide. A fully biologically active molecule, rhGM-CSF has a molecular mass of 26-32kDa analyzed by SDS-PAGE and is obtained by proprietary refolding and chromatographic techniques at GenScript.

Amino Acid Sequence:

```
00001 MAPARSPSPS TQPWEHVNAI QEALRLLNLS RDAAEMNET
00041 VEISEMFDL QEPTCLQTRL ELYKQGLRGS LTKLKGPLTM
00081 MASHYKQHCP PTPETSCATQ IITFESFKEN LKDFLLVIPF
00121 DCWEPVQE
```

Source: *P. pastoris*

Species: Human

Biological Activity: ED₅₀ < 0.2ng/ml, measured by proliferation assay of TF-1 cells, corresponding to a specific activity of > 5x 10⁶ units/mg.

Molecular Weight: 26-32kDa, observed by SDS-PAGE.

Formulation: Lyophilized after extensive dialysis against 10 mM PB, pH7.0.

Reconstitution: Reconstituted in ddH₂O at 100 µg/ml.

Purity: > 95 % as analyzed by SDS-PAGE.

Endotoxin Level: <1.0EU/µg, determined by LAL method.

Storage: Lyophilized recombinant human Granulocyte Macrophage Colony Stimulating Factor (rhGM-CSF) remains stable up to 12 months at lower than -70°C from date of receipt. Upon reconstitution, rhGM-CSF should be stable up to 4 weeks at 4°C or up to 6 months at -20°C.