

## IL-11, Mouse

**Cat. No.:** Z02769-10

**Size:** 10.0 ug

**Synonyms:** Interleukin-11 (IL-11), Mouse;

### Description:

Interleukin 11 is a pleiotropic cytokine that was originally detected in the conditioned medium of an IL-1<sup>a</sup>-stimulated primate bone marrow stromal cell line (PU-34) as a mitogen for the IL-6-responsive mouse plasmacytoma cell line T11. IL-11 contains no cysteine residues or potential glycosylation sites. IL-11 has multiple effects on both hematopoietic and non-hematopoietic cells. Many of the biological effects described for IL-11 overlap those for IL-6. In vitro, IL-11 can synergize with IL-3, IL-4 and SCF to shorten the G0 period of early hematopoietic progenitors. IL-11 also enhances the IL-3-dependent megakaryocyte colony formation. IL-11 has been found to stimulate the T cell dependent development of specific immunoglobulin-secreting B cell.

### Amino Acid Sequence:

```
00001 MPGPPAGSPR VSSDPRADLD SAVLLTRSLL ADTRQLAAQM
00041 RDKFPADGDH SLDSLPTLAM SAGTLGSLQL PGVLRRLRVD
00081 LMSYLRHVQW LRRAGGPSLK TLEPELGALQ ARLERLLRRL
00121 QLLMSRLALP QAAPDQVIP LGPPASAWGS IRAAHAILGG
00161 LHLLTDWAVR GLLLLKTRL
```

**Source:** *E. coli*

**Species:** Mouse

**Biological Activity:** Fully biologically active when compared to standard. The ED<sub>50</sub> as determined by a cell proliferation assay using murine T11 cells is less than 2 ng/ml, corresponding to a specific activity of > 5.0 × 10<sup>5</sup> IU/mg.

**Molecular Weight:** Approximately 19.1 kDa, a single non-glycosylated polypeptide chain containing 179 amino acids.

**Formulation:** Lyophilized from a 0.2 μm filtered solution in PBS, pH 7.4.

**Appearance:** Sterile Filtered White lyophilized (freeze-dried) powder.

**Reconstitution:** We recommend that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. Reconstitute in sterile distilled water or aqueous buffer containing 0.1 % BSA to a concentration of 0.1-1.0 mg/mL. Stock solutions should be apportioned into working aliquots and stored at ≤ -20 °C. Further dilutions should be made in appropriate buffered solutions.

**Purity:** > 97 % by SDS-PAGE and HPLC analyses.

**Endotoxin Level:** Less than 1 EU/μg of rMuIL-11 as determined by LAL method.

**Storage:** This lyophilized preparation is stable at 2-8 °C, but should be kept at -20 °C for long term storage, preferably desiccated. Upon reconstitution, the preparation is stable for up to one week at 2-8 °C. For maximal stability, apportion the reconstituted preparation into working aliquots and store at -20 °C to -70 °C. Avoid repeated freeze/thaw cycles.