

I-TAC/CXCL11, Human

Cat. No.: Z02824-20

Size: 20.0 ug

Synonyms: I-TAC/CXCL11, Human;

Description:

CXCL11 cDNA encodes a 94 amino acid (aa) residue precursor protein with a 21 aa residue putative signal sequence, which is cleaved to form the mature 73 aa residue protein. CXCL11 shares 36% and 37% amino acid sequence homology with IP-10 and MIG (two other known human non-ELR CXC chemokines), respectively. CXCL11 is expressed at low levels in normal tissues including thymus, spleen and pancreas. The expression of CXCL11 mRNA is radically up regulated in IFN- γ and IL-1 stimulated astrocytes. Moderate increase in expression is also observed in stimulated monocytes. CXCL11 has potent chemoattractant activity for IL-2 activated T cells and transfected cell lines expressing CXCR3, but not freshly isolated T-cells, neutrophils or monocytes.

Amino Acid Sequence:

00001 FPMFKRGRCL CIGPGVKAVK VADIEKASIM YPSNCDKIE
00041 VIITLKENKG QRCLNPKSKQ ARLIKKVER KNF

Source: *E. coli*

Species: Human

Biological Activity: Fully biologically active when compared to standard. The biological activity determined by a chemotaxis bioassay using human IL-2 activated human T-lymphocytes is in a concentration range of 0.1-10 ng/ml.

Molecular Weight: Approximately 8.3 kDa, a single non-glycosylated polypeptide chain containing 73 amino acids.

Formulation: Lyophilized from a 0.2 μ m filtered concentrated solution in 20 mM PB, pH 7.4, 100 mM NaCl.

Appearance: Sterile Filtered White lyophilized (freeze-dried) powder.

Reconstitution: We recommend that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. Reconstitute in sterile distilled water or aqueous buffer containing 0.1 % BSA to a concentration of 0.1-1.0 mg/mL. Stock solutions should be apportioned into working aliquots and stored at ≤ -20 °C. Further dilutions should be made in appropriate buffered solutions.

Purity: > 97 % by SDS-PAGE and HPLC analyses.

Endotoxin Level: Less than 1 EU/ μ g of rHu-I-TAC/CXCL11 as determined by LAL method.

Storage: This lyophilized preparation is stable at 2-8 °C, but should be kept at -20 °C for long term storage, preferably desiccated. Upon reconstitution, the preparation is stable for up to one week at 2-8 °C. For maximal stability, apportion the reconstituted preparation into working aliquots and store at -20 °C to -70 °C. Avoid repeated freeze/thaw cycles.