

OSM, Rat

Cat. No.: Z03144-50

Size: 50.0 ug

Synonyms: OSM

Description:

Oncostatin M (OSM) is a multifunctional cytokine, and belongs to Interleukin-6 (IL-6) subfamily, which also includes IL-11, leukemia inhibitory factor (LIF), ciliary neurotropic factor, cardiotrophin-1, and novel neurotrophin-1. *In vivo*, OSM is secreted from activated T cells, monocytes, neutrophils, and endothelial cells. OSM is related to LIF, and shares a receptor with LIF in human. Human OSM can bind to gp130 and recruit OSM Receptor β or LIF Receptor β to form a ternary complex. OSM stimulates the growth of different types of cells, including megakaryocytes, fibroblasts, vascular endothelial cells, and T cells. OSM inhibits the proliferation of several cancer cell lines, such as solid tissue tumor cells, lung cancer cells, melanoma cells, and breast cancer cells. Recombinant Rat Oncostatin M (rrOSM) produced in *E. coli* is a single non-glycosylated polypeptide chain containing 215 amino acids. A fully biologically active molecule, rrOSM has a molecular mass of 24.5 kDa analyzed by reducing SDS-PAGE and is obtained by proprietary chromatographic techniques at GenScript.

Amino Acid Sequence:

```
00001 MKRGCSSSP KLLSQLKSQA NITGNTASLL EPYILHQLN  
00041 TLTLRAACTE HPVAFPSDEM LRQLSKPDFL STVHATLGRV  
00081 WHQLGAFRQQ FPKIQDFPEL ERARQNIQGI RNNVYCMARL  
00121 LHPPLEIPEP TQADSGTSRP TTTAPGIFQI KIDSCRFLWG  
00161 YHRFMGSGVR VFEWGDGSR RSRRHSPWA WLKGDHRIRP  
00201 SRSSQSAMLR SLVPR
```

Source: *E. coli*

Species: Rat

Biological Activity: ED₅₀ < 10 ng/mL, measured by a cell proliferation assay using Rat Embryo Brain cells, corresponding to a specific activity of > 1 × 10⁵ units/mg.

Molecular Weight: 24.5 kDa, observed by reducing SDS-PAGE.

Formulation: Lyophilized after extensive dialysis against PBS.

Reconstitution: Reconstituted in ddH₂O or PBS at 100 µg/ml.

Purity: > 95% as analyzed by SDS-PAGE and HPLC.

Endotoxin Level: < 0.2 EU/µg, determined by LAL method.

Storage: Lyophilized recombinant Rat Oncostatin M (rrOSM) remains stable up to 6 months at lower than -70°C from date of receipt. Upon reconstitution, rrOSM remains stable up to 2 weeks at 4°C or up to 3 months at -20°C.