

Rev03
Update: Dec,14,2021

DATASHEET

PEDF, Human

Cat. No.: Z02722

Product Introduction

Species	Human
Protein Construction	PEDF (Gln20-Pro418) Accession # P36955
Purity	> 97% as analyzed by SDS-PAGE > 97% as analyzed by HPLC
Endotoxin Level	< 1 EU/ μ g of protein by LAL method
Biological Activity	Fully biologically active when compared to standard. The ED ₅₀ as determined by its ability to enhance the adhesion of human Saos2 cells to bovine Collagen I coated plate is less than 2.0 ng/ml, corresponding to a specific activity of $> 5.0 \times 10^5$ IU/mg.
Expression System	E. coli
Theoretical Molecular Weight	44.4 kDa
Formulation	Lyophilized from a 0.2 μ m filtered solution in 20 mM PB, pH 7.4, 150 mM NaCl.
Reconstitution	It is recommended that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. Reconstitute the lyophilized powder in sterile distilled water or aqueous buffer containing 0.1 % BSA to a concentration of 0.1-1.0 mg/ml.
Storage & Stability	Upon receiving, this product remains stable for up to 6 months at -70°C or -20°C. Upon reconstitution, the product should be stable for up to 1 week at 4°C or up to 3 months at -20°C. Avoid repeated freeze-thaw cycles.

Background

Target Background : PEDF is a noninhibitory serpin with neurotrophic, anti-angiogenic, and anti-tumorigenic properties. It is a 50 kDa glycoprotein produced and secreted in many tissues throughout the body. A major component of the anti-angiogenic action of PEDF is the induction of apoptosis in proliferating endothelial cells. In addition, PEDF is able to inhibit the activity of angiogenic factors such as VEGF and FGF-2. The neuroprotective effects of PEDF are achieved through suppression of neuronal apoptosis induced by peroxide, glutamate, or other neurotoxins. The recent identification of a lipase-linked cell membrane receptor for PEDF (PEDF-R) that binds to PEDF with high affinity should facilitate further elucidation of the underlying mechanisms of this pluripotent serpin. To date, PEDF-R is the only signaling receptor known to be used by a serpin family member. The unique range of PEDF activities implicate it as a potential therapeutic agent for the treatment of vasculature related neurodegenerative diseases such as age-related macular degeneration (AMD) and proliferative diabetic retinopathy (PDR). PEDF also has the potential to be useful in the treatment of various angiogenesis-related diseases including a number of cancers.

Synonyms : SERPINF1; EPC-1; OI12; OI6; PEDF; PIG35; serpin family F member 1; Pigment epithelium-derived factor

For laboratory research use only. Direct human use, including taking orally and injection and clinical use are forbidden.