

Rev04  
 Update: Mar,01,2022

## DATASHEET

# Flt-3L, RhesusMacaque

Cat. No.: Z02762

## Product Introduction

<b>Species</b>	Rhesus Macaque
<b>Protein Construction</b>	Flt-3L (Thr27-Pro185) Accession # H9Z6V7
<b>Purity</b>	> 97% as analyzed by SDS-PAGE > 97% as analyzed by HPLC
<b>Endotoxin Level</b>	< 1 EU/μg of protein by LAL method
<b>Biological Activity</b>	Fully biologically active when compared to standard. The ED <sub>50</sub> as determined by a cell proliferation assay using human AML5 cells is less than 1.0 ng/ml, corresponding to a specific activity of > 1.0 × 10 <sup>6</sup> IU/mg.
<b>Expression System</b>	E. coli
<b>Theoretical Molecular Weight</b>	18 kDa
<b>Formulation</b>	Lyophilized from a 0.2 μm filtered solution in PBS, pH 7.4.
<b>Reconstitution</b>	It is recommended that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. Reconstitute the lyophilized powder in sterile distilled water or aqueous buffer containing 0.1% BSA to a concentration of 0.1-1.0 mg/ml.
<b>Storage &amp; Stability</b>	Upon receiving, this product remains stable for up to 6 months at -70°C or -20°C. Upon reconstitution, the product should be stable for up to 1 week at 4°C or up to 3 months at -20°C. Avoid repeated freeze-thaw cycles.

## Background

**Target Background :** Flt-3 ligand (FL) is a recently identified hematopoietic cytokine whose activities are mediated by binding to the transmembrane glycoprotein Flt-3. Flt-3 was first discovered as a member of the class III subfamily of receptor tyrosine kinases (RTK) whose expression among hematopoietic cells was found to be restricted to highly enriched stem/progenitor cell populations. Additional class III RTKs include the receptors from SCF, M-CSF and PDGF. Not surprisingly, Flt-3 ligand is also structurally related to M-CSF and SCF. All three cytokines have been shown to exist both as type I transmembrane proteins and as soluble proteins.

**Synonyms :** Flt3 ligand; Flt3LG; SL cytokine; Fms-related tyrosine kinase 3 ligand

For laboratory research use only. Direct human use, including taking orally and injection and clinical use are forbidden.