

Rev03
Update: Dec,14,2021

DATASHEET

GRO- γ /CXCL3, Human

Cat. No.: Z02819

Product Introduction

Species	Human
Protein Construction	GRO-γ (Ala35-Asn107) Accession # P19876
Purity	> 97% as analyzed by SDS-PAGE > 97% as analyzed by HPLC
Endotoxin Level	< 1 EU/ μ g of protein by LAL method
Biological Activity	Fully biologically active when compared to standard. The biological activity determined by a chemotaxis bioassay using human CXCR2 transfected human 293 cells is in a concentration range of 10.0-100.0 ng/ml.
Expression System	E. coli
Theoretical Molecular Weight	7.9 kDa
Formulation	Lyophilized from a 0.2 μ m filtered solution in 20 mM PB, pH 7.4, 50 mM NaCl.
Reconstitution	It is recommended that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. Reconstitute the lyophilized powder in sterile distilled water or aqueous buffer containing 0.1% BSA to a concentration of 0.1-1.0 mg/ml.
Storage & Stability	Upon receiving, this product remains stable for up to 6 months at -70°C or -20°C. Upon reconstitution, the product should be stable for up to 1 week at 4°C or up to 3 months at -20°C. Avoid repeated freeze-thaw cycles.

Background

Target Background : Chemokine (C-X-C motif) ligand 3 (CXCL3) is a small cytokine belonging to the CXC chemokine family that is also known as GRO3 oncogene (GRO3), GRO protein gamma (GRO γ) and macrophage inflammatory protein-2-beta (MIP2b). CXCL3 controls migration and adhesion of monocytes and mediates its effect on its target cell by interacting with cell surface chemokine receptor CXCR2. It has been shown that CXCL3 regulates the migration of precursors of cerebellar granule neurons toward the internal layers of the cerebellum, during morphogenesis.

Synonyms : GRO-gamma; CXCL3

For laboratory research use only. Direct human use, including taking orally and injection and clinical use are forbidden.