

Rev03
Update: Dec,14,2021

DATASHEET

NAP-2/CXCL7, Rat

Cat. No.: Z02956

Product Introduction

Species	Rat
Protein Construction	CXCL7 (Ile46-Ile107) Accession # Q99ME0
Purity	> 97% as analyzed by SDS-PAGE > 97% as analyzed by HPLC
Endotoxin Level	< 1 EU/ μ g of protein by LAL method
Biological Activity	Fully biologically active when compared to standard. The biologically active determined by a chemotaxis bioassay using human CXCR2 transfected murine BaF3 cells is in a concentration range of 0.1-1.0 ng/ml.
Expression System	E. coli
Theoretical Molecular Weight	6.8 kDa
Formulation	Lyophilized from a 0.2 μ m filtered solution in PBS, pH 7.4.
Reconstitution	It is recommended that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. Reconstitute the lyophilized powder in sterile distilled water or aqueous buffer containing 0.1 % BSA to a concentration of 0.1-1.0 mg/ml.
Storage & Stability	Upon receiving, this product remains stable for up to 6 months at -70°C or -20°C. Upon reconstitution, the product should be stable for up to 1 week at 4°C or up to 3 months at -20°C. Avoid repeated freeze-thaw cycles.

Background

Target Background : Neutrophil Activating Peptide 2 (NAP-2) is proteolytically processed carboxyl-terminal fragments of platelet basic protein (PBP) which is found in the alpha-granules of human platelets. NAP-2 is a member of the CXC chemokines. Similar to other ELR domain containing CXC chemokines such as IL-8 and the GRO proteins, NAP-2 has been shown to bind CXCR-2 and to chemoattract and activate neutrophils. Although CTAP-III, β -TG and PBP represent amino-terminal extended variants of NAP-2 and possess the same CXC chemokine domains, these proteins do not exhibit NAP-2 activity. Recently, it has been shown that the additional amino-terminal residues of CTAP-III masks the critical ELR receptor binding domain that is exposed on NAP-2 and may account for lack of NAP-2 activity.

Synonyms : PPBP; B-TG1; Beta-TG; CTAP-III; CTAP3; CTAPIII; CXCL-7; LA-PF4; LDGF; MDGF; NAP2; PBP; SCYB7; TC1; TC2; TGB; TGB1; THBGB; THBGB1; pro-platelet basic protein

For laboratory research use only. Direct human use, including taking orally and injection and clinical use are forbidden.