

Rev03  
Update: Dec,14,2021

**DATASHEET**

# CXCL17, Rat

Cat. No.: Z02957

## Product Introduction

<b>Species</b>	Rat
<b>Protein Construction</b>	<b>CXCL17 (Ser23-Leu119)</b> Accession # D4A875
<b>Purity</b>	> 97% as analyzed by SDS-PAGE > 97% as analyzed by HPLC
<b>Endotoxin Level</b>	< 1 EU/ $\mu$ g of protein by LAL method
<b>Biological Activity</b>	Fully biologically active when compared to standard. The ED <sub>50</sub> as determined by its ability to induce VEGF expression using murine endothelial cells is less than 5.0 $\mu$ g/ml, corresponding to a specific activity of > 200 IU/mg.
<b>Expression System</b>	E. coli
<b>Theoretical Molecular Weight</b>	11.5 kDa
<b>Formulation</b>	Lyophilized from a 0.2 $\mu$ m filtered solution in 2 $\times$ PBS, pH 7.4.
<b>Reconstitution</b>	It is recommended that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. Reconstitute the lyophilized powder in sterile distilled water or aqueous buffer containing 0.1% BSA to a concentration of 0.1-1.0 mg/ml.
<b>Storage &amp; Stability</b>	Upon receiving, this product remains stable for up to 6 months at -70°C or -20°C. Upon reconstitution, the product should be stable for up to 1 week at 4°C or up to 3 months at -20°C. Avoid repeated freeze-thaw cycles.

## Background

**Target Background** : Chemokine (C-X-C motif) ligand 17 (CXCL17) is a small cytokine belonging to the CXC chemokine family that has been identified in humans and mice. CXCL17 attracts dendritic cells and monocytes and is regulated in tumors. It is also known as VEGF co-regulated chemokine 1 (VCC-1) and dendritic cell- and monocyte-attracting chemokine-like protein (DMC). This chemokine is constitutively expressed in the lung.

**Synonyms** : Protein Cxcl17; Cxcl17; RGD1304717; C-X-C motif chemokine 17; VEGF co-regulated chemokine 1; Vcc1; VCC-1

For laboratory research use only. Direct human use, including taking orally and injection and clinical use are forbidden.