

Rev04
Update: Jan,19,2024

DATASHEET


IL-4, Mouse

Cat. No.: Z02996


Product Introduction

Species	Mouse
Protein Construction	IL-4 (His21-Ser140) Accession # P07750
Purity	> 95% as analyzed by SDS-PAGE > 95% as analyzed by HPLC
Endotoxin Level	< 0.2 EU/μg of protein by gel clotting method
Biological Activity	ED ₅₀ < 2.0 ng/ml, measured in a cell proliferation assay using murine HT-2 cells, corresponding to a specific activity of > 5.0 × 10 ⁵ units/mg.
Expression System	CHO
Apparent Molecular Weight	~15 kDa, on SDS-PAGE under reducing conditions.
Formulation	Lyophilized after extensive dialysis against PBS.
Reconstitution	It is recommended that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. Reconstitute the lyophilized powder in ddH ₂ O or PBS up to 100 μg/ml.
Storage & Stability	Upon receiving, this product remains stable for up to 6 months at lower than -70°C. Upon reconstitution, the product should be stable for up to 1 week at 4°C or up to 3 months at -20°C. For long term storage it is recommended that a carrier protein (example 0.1% BSA) be added. Avoid repeated freeze-thaw cycles.

Examples



ED₅₀ < 2.0 ng/ml, measured in a cell proliferation assay using murine HT-2 cells, corresponding to a specific activity of > 5.0 × 10⁵ units/mg.



Lane 1: 2 µg of IL-4, Mouse, non-reducing (NR)
 Lane 2: 2 µg of IL-4, Mouse, reducing (R)
 > 95% as analyzed by SDS-PAGE

Background

Target Background : Interleukin-4 (IL-4) is a pleiotropic cytokine regulates diverse T and B cell responses including cell proliferation, survival, and gene expression. It has important effects on the growth and differentiation of different immunologically competent cells. Interleukin-4 is produced by mast cells, T cells, and bone marrow stromal cells. IL-4 regulates the differentiation of native CD4⁺ T cells (Th0 cells) into helper Th2 cells, and regulates the immunoglobulin class switching to the IgG1 and IgE isotypes. IL-4 has numerous important biological functions including stimulating B-cells activation, T-cell proliferation and CD4⁺ T-cells differentiation to Th2 cells. It is a key regulator in hormone control and adaptive immunity. IL-4 also plays a major role in inflammation response and wound repair via activation of macrophage into M2 cells. IL-4 is stabilized by three disulphide bonds forming a compact globular protein structure. Four alpha-helix bundle with left-handed twist is dominated half of the protein structure with 2 overhand connections and fall into a 2-stranded anti-parallel beta sheet.

Synonyms : IL4; B cell growth factor-1; BCGF-1; B cell stimulatory factor-1; BSF-1; B cell differentiation factor; BCDF; T cell growth factor-2; TCGF-2; Mast-cell growth factor-2; MCGF-2

For laboratory research use only. Direct human use, including taking orally and injection and clinical use are forbidden.

生产商：南京金斯瑞生物科技有限公司 江苏省南京市江宁区科学园雍熙路28号

Manufacturer: Nanjing GenScript Biotech Co., Ltd. No. 28Yongxi Road, Jiangning District, Nanjing, Jiangsu, China