

Rev03  
Update: Dec,14,2021

**DATASHEET**

# AITRL, Mouse

Cat. No.: Z03098

## Product Introduction

<b>Species</b>	Mouse
<b>Protein Construction</b>	<b>AITRL (Thr47-Ser173) Accession # Q7TS55</b>
<b>Purity</b>	> 95% as analyzed by SDS-PAGE > 95% as analyzed by HPLC
<b>Endotoxin Level</b>	< 0.2 EU/μg of protein by gel clotting method
<b>Biological Activity</b>	ED <sub>50</sub> < 5.0 ng/ml, measured by the amount of Interleukin-8 produced by PMBC, corresponding to a specific activity of > 2.0 × 10 <sup>5</sup> units/mg.
<b>Expression System</b>	E. coli
<b>Apparent Molecular Weight</b>	~14.5 kDa, on SDS-PAGE under reducing conditions.
<b>Formulation</b>	Lyophilized after extensive dialysis against PBS.
<b>Reconstitution</b>	It is recommended that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. Reconstitute the lyophilized powder in ddH <sub>2</sub> O or PBS up to 100 μg/ml.
<b>Storage &amp; Stability</b>	Upon receiving, this product remains stable for up to 6 months at lower than -70°C. Upon reconstitution, the product should be stable for up to 1 week at 4°C or up to 3 months at -20°C. For long term storage it is recommended that a carrier protein (example 0.1% BSA) be added. Avoid repeated freeze-thaw cycles.

## Background

**Target Background** : Activation-Inducible TNF-Related Ligand (AITRL), also known as Glucocorticoid-Induced TNF-Related Ligand (GITRL), belongs to the tumor necrosis factor superfamily (TNFSF). AITRL is a Type II single transmembrane protein and shares low conservation within the extracellular domain with other TNFSF members. AITRL is expressed on macrophages, immature and mature dendritic cells and B cells. Its receptor, Activation-Inducible TNFR family Receptor (AITR), is expressed on T lymphocytes, natural killer (NK) cells, and antigen-presenting cells. After binding by AITRL, AITR can be released. AITR activation increases resistance to tumors and viral infections and is involved in autoimmune and inflammatory processes. In addition, activated AITR increases TCR-induced T cell proliferation and cytokine production and rescues T cells and NK cells from apoptosis.

**Synonyms** : Activation-induced TNFR member Ligand; TNFSF18; GITRL; TL-6

**GenScript USA, Inc.**

860 Centennial Ave. Piscataway, NJ 08854

Tel: 1-732-885-9188

For laboratory research use only. Direct human use, including taking orally and injection and clinical use are forbidden.