

Rev03 DATASHEET

Update: Dec,14,2021

FGF-18, Rat

Cat. No.: Z03100

Product Introduction

Species	Rat
Protein Construction	FGF-18 (Glu28-Arg199) Accession # 088182
Purity	> 95% as analyzed by SDS-PAGE > 95% as analyzed by HPLC
Endotoxin Level	< 0.2 EU/µg of protein by gel clotting method
Biological Activity	ED $_{50}$ < 0.5 µg/ml, measured by a cell proliferation assay using 3T3 cells, corresponding to a specific activity of > 2.0 \times 10 ³ units/mg.
Expression System	E. coli
Apparent Molecular Weight	~20.1 kDa, on SDS-PAGE under non-reducing conditions.
Formulation	Lyophilized after extensive dialysis against PBS.
Reconstitution	It is recommended that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. Reconstitute the lyophilized powder in ddH ₂ O up to 100 μg/ml.
Storage & Stability	Upon receiving, this product remains stable for up to 6 months at lower than -70°C. Upon reconstitution, the product should be stable for up to 1 week at 4°C or up to 3 months at -20°C. For long term storage it is recommended that a carrier protein (example 0.1% BSA) be added. Avoid repeated freeze-thaw cycles.

Background

Target Background: Fibroblast Growth Factor 18 (FGF-18) is a pleiotropic cytokine belonging to the heparin-binding FGF family, which has 23 different members. Structurally, FGF-18 is closely related to FGF-8 and FGF-17. Like other FGFs, FGF-18 can bind to different FGF receptors in vivo. FGF-18 is expressed in various tissues and has multiple functions: during long bone growth, FGF-18 is expressed in perichondrium and developing joints, and regulates bone formation by inhibiting chondrocyte proliferation and differentiation; FGF-18 knock-out mice survive embryonic development, but exhibit skeletal abnormalities and die in the early neonatal period. FGF-18 also induces ectopic cartilage formation in the lung, and alters the morphology of the pulmonary mesenchyma.

Synonyms: FGF18; Fibroblast Growth Factor-18; zFGF5; FGFI



For laboratory research use only. Direct human use, including taking orally and injection and clir orbidden.	nical use are