

Rev04
Update: Mar,01,2022

DATASHEET

Betacellulin, Human

Cat. No.: Z03102

Product Introduction

Species	Human
Protein Construction	Betacellulin (Asp32-Tyr111) Accession # P35070
Purity	> 95% as analyzed by SDS-PAGE
Endotoxin Level	< 0.2 EU/ μ g of protein by gel clotting method
Biological Activity	ED ₅₀ < 4.0 pg/ml, measured in a cell proliferation assay using 3T3 cells.
Expression System	HEK 293
Apparent Molecular Weight	15~18 kDa, on SDS-PAGE under reducing conditions.
Formulation	Lyophilized after extensive dialysis against PBS.
Reconstitution	It is recommended that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. Reconstitute the lyophilized powder in ddH ₂ O or PBS up to 100 μ g/ml.
Storage & Stability	Upon receiving, this product remains stable for up to 6 months at lower than -70°C. Upon reconstitution, the product should be stable for up to 1 week at 4°C or up to 3 months at -20°C. For long term storage it is recommended that a carrier protein (example 0.1% BSA) be added. Avoid repeated freeze-thaw cycles.

Background

Target Background : Betacellulin (BTC) is a member of the EGF family of growth factors that also includes EGF, TGF- α , Amphiregulin, HB-EGF, Epiregulin, Tomoregulin, Heregulin and Neuregulins. Mature human BTC protein exhibits 80% amino acidsimilarity with mouse BTC protein. BTC is expressed in most tissues including kidney, uterus, liver and pancreas. It is also present in body fluids, including serum, milk, and colostrum. It is synthesized primarily as a transmembrane precursor, which is then processed to a mature molecule by proteolytic events. BTC signals through the EGF receptor.

Synonyms : BTC

For laboratory research use only. Direct human use, including taking orally and injection and clinical use are forbidden.