

Rev03
 Update: Dec,14,2021

DATASHEET

IL-7, His, Mouse


Cat. No.: Z03209

Product Introduction

Species	Mouse
Protein Construction	<div style="display: flex; align-items: center; gap: 10px;"> <div style="background-color: #0056b3; color: white; padding: 5px; text-align: center;"> IL-7 (Glu26-Ile154) Accession # Q544C8 </div> <div style="background-color: #76b82a; color: white; padding: 5px; text-align: center;"> Poly-His </div> </div> <div style="display: flex; justify-content: space-around; margin-top: 5px; font-size: small;"> N-term C-term </div>
Purity	> 95% as analyzed by SDS-PAGE
Endotoxin Level	< 0.2 EU/μg of protein by gel clotting method
Biological Activity	ED ₅₀ < 0.4 ng/ml, measured in a cell proliferation assay using 2E8 cells.
Expression System	CHO
Apparent Molecular Weight	8~28kDa, on SDS-PAGE under reducing conditions.
Formulation	Lyophilized after extensive dialysis against PBS.
Reconstitution	It is recommended that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. Reconstitute the lyophilized powder in ddH ₂ O or PBS up to 100 μg/ml.
Storage & Stability	Upon receiving, this product remains stable for up to 6 months at lower than -70°C. Upon reconstitution, the product should be stable for up to 1 week at 4°C or up to 3 months at -20°C. For long term storage it is recommended that a carrier protein (example 0.1% BSA) be added. Avoid repeated freeze-thaw cycles.

Examples

ED₅₀<0.4ng/ml, measured in a cell proliferation assay using 2E8 cells.



Background

Target Background : Interleukin-7 (IL-7), also known as lymphopoietin 1 and pre-B cell factor, is a hematopoietic growth factor belonging to the IL-7/IL-9 family. It is produced by keratinocytes, dendritic cells, hepatocytes, neurons and epithelial cells. IL-7 binds and signals through IL-7 receptor, a heterodimer consisting of IL-7 receptor alpha and common gamma chain receptor. IL-7 plays a role in regulating early B cell and T cell development. It is also important for optimal dendritic cell-T cell interaction.

Synonyms : IL7; IL-7; interleukin 7; Lymphopoietin 1; LP-1; pre-B cell factor

For laboratory research use only. Direct human use, including taking orally and injection and clinical use are forbidden.