

Rev04
Update: Mar,01,2022

DATASHEET

MPIF-1/CCL23, Human

Cat. No.: Z03248

Product Introduction

Species	Human
Protein Construction	MPIF-1 (Arg22-Asn120) Accession # P55773
Purity	> 98% as analyzed by SDS-PAGE
Endotoxin Level	< 0.2 EU/μg of protein by gel clotting method
Biological Activity	The EC ₅₀ value of human MPIF-1/CCL23 on Ca ²⁺ mobilization assay in CHO-K1/Ga15/hCCR1 cells (human Ga15 and human CCR1 stably expressed in CHO-K1 cells) is less than 2.0 μg/ml.
Expression System	CHO
Apparent Molecular Weight	~12 kDa, on SDS-PAGE under reducing conditions.
Formulation	Lyophilized after extensive dialysis against PBS.
Reconstitution	It is recommended that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. Reconstitute the lyophilized powder in ddH ₂ O or PBS up to 100 μg/ml.
Storage & Stability	Upon receiving, this product remains stable for up to 6 months at lower than -70°C. Upon reconstitution, the product should be stable for up to 1 week at 4°C or up to 3 months at -20°C. For long term storage it is recommended that a carrier protein (example 0.1% BSA) be added. Avoid repeated freeze-thaw cycles.

Background

Target Background : Myeloid progenitor inhibitory factor 1 (MPIF-1), also known as Chemokine (C-C motif) ligand 23 (CCL23) is a small cytokine belonging to the CC chemokine family. MPIF-1 is predominantly expressed in lung and liver tissue, but is also found in bone marrow and placenta. It is also expressed in some cell lines of myeloid origin. It is highly chemotactic for resting T cells and monocytes and slightly chemotactic for neutrophils. MPIF-1 has been shown to inhibit colony formation of bone marrow myeloid immature progenitors. It has also been attributed to an inhibitory activity on hematopoietic progenitor cells. MPIF-1 is a ligand for the chemokine receptor CCR1.

Synonyms : SCYA23; Ckb-8; MPIF-1; MIP-3; CKb8; Macrophage Inflammatory Protein-3; CCL-23; Ck-beta8

For laboratory research use only. Direct human use, including taking orally and injection and clinical use are forbidden.