

Rev03
Update: Dec,14,2021

DATASHEET

DCIP-1/CXCL3, Mouse

Cat. No.: Z03249

Product Introduction

Species	Mouse
Protein Construction	CXCL3 (Ala28-Ser100) Accession # Q6W5C0
Purity	> 98% as analyzed by SDS-PAGE
Endotoxin Level	< 0.2 EU/μg of protein by gel clotting method
Biological Activity	The EC ₅₀ value of mouse DCIP-1/CXCL3 on Ca ²⁺ mobilization assay in CHO-K1/Ga15/mCXCR2 cells (human Ga15 and mouse CXCR2 stably expressed in CHO-K1 cells) is less than 100.0 ng/ml.
Expression System	CHO
Apparent Molecular Weight	~8 kDa, on SDS-PAGE under reducing conditions.
Formulation	Lyophilized after extensive dialysis against PBS.
Reconstitution	It is recommended that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. Reconstitute the lyophilized powder in ddH ₂ O or PBS up to 100 μg/ml.
Storage & Stability	Upon receiving, this product remains stable for up to 6 months at lower than -70°C. Upon reconstitution, the product should be stable for up to 1 week at 4°C or up to 3 months at -20°C. For long term storage it is recommended that a carrier protein (example 0.1% BSA) be added. Avoid repeated freeze-thaw cycles.

Background

Target Background : Chemokine (C-X-C motif) ligand 3 (CXCL3) is a small cytokine belonging to the CXC chemokine family that is also known as DCIP-1 (dendritic cell inflammatory protein-1) or MIP2b. CXCL3 controls migration and adhesion of monocytes and mediates its effect on its target cell by interacting with cell surface chemokine receptor CXCR2. It has been shown that CXCL3 regulates the migration of precursors of cerebellar granule neurons toward the internal layers of cerebellum, during morphogenesis.

Synonyms : C-X-C motif chemokine 3; Dendritic cell inflammatory protein 1; Dcip1; Gm1960

For laboratory research use only. Direct human use, including taking orally and injection and clinical use are forbidden.