

Rev04
Update: Feb,23,2022

DATASHEET

LIX/CXCL5 (74aa), Mouse

Cat. No.: Z03250

Product Introduction

Species	Mouse
Protein Construction	CXCL5 (Val45-Ala118) Accession # P50228
Purity	> 98% as analyzed by SDS-PAGE
Endotoxin Level	< 0.2 EU/μg of protein by gel clotting method
Biological Activity	The EC ₅₀ value of mouse LIX/CXCL5 (74aa) on Ca ²⁺ mobilization assay in CHO-K1/Gα15/mCXCR2 cells (human Gα15 and mouse CXCR2 stably expressed in CHO-K1 cells) is less than 200.0 ng/ml.
Expression System	HEK 293 Cells
Apparent Molecular Weight	~8 kDa, on SDS-PAGE under reducing conditions.
Formulation	Lyophilized after extensive dialysis against PBS.
Reconstitution	It is recommended that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. Reconstitute the lyophilized powder in ddH ₂ O or PBS up to 100 μg/ml.
Storage & Stability	Upon receiving, this product remains stable for up to 6 months at lower than -70°C. Upon reconstitution, the product should be stable for up to 1 week at 4°C or up to 3 months at -20°C. For long term storage it is recommended that a carrier protein (example 0.1% BSA) be added. Avoid repeated freeze-thaw cycles.

Background

Target Background : Mouse LIX (C-X-C motif chemokine 5) is a small cytokine belonging to the CXC chemokine family that is cleaved into the following 2 chains [GCP-2(1-78) and GCP-2(9-78)]. Mouse LIX plays a role in reducing sensitivity to sunburn pain in some subjects, and is a potential target which could be used to understand more about pain in other inflammatory conditions. It is most closely related to two highly homologous human neutrophil chemoattractants GCP-2 and ENA-78. The first 78 amino acid residues within the predicted mature mouse LIX shares approximately 61% and 55% amino acid identity with human GCP-2 and ENA-78. This chemokine stimulates the chemotaxis of neutrophils possessing angiogenic properties. It elicits these effects by interacting with the cell surface chemokine receptor CXCR2.

Synonyms : C-X-C motif chemokine 5; Small-inducible cytokine B5; Cytokine LIX; Scyb5; GCP-2; Scyb6; ENA-78; AMCF-II; Epithelial Neutrophil Activating Peptide-78

For laboratory research use only. Direct human use, including taking orally and injection and clinical use are forbidden.