

Rev03 DATASHEET

Update: Dec,14,2021

I-309/CCL1, Human

Cat. No.: Z03273

Product Introduction

| Species | Human |
|------------------------------|--|
| Protein Construction | CCL1 (Ser23-Lys96) Accession # P22362-1 |
| Purity | > 95% as analyzed by SDS-PAGE |
| Endotoxin Level | < 0.2 EU/µg of protein by gel clotting method |
| Biological Activity | The EC $_{50}$ value of human I-309/CCL on Ca $^{2+}$ mobilization assay in CHO-K1/G α 15/hCCR8 cells (human G α 15 and human CCR8 stably expressed in CHO-K1 cells) is less than 1.0 μ g/ml. |
| Expression System | E. coli |
| Theoretical Molecular Weight | 8.6 kDa |
| Apparent Molecular Weight | ~8.5 kDa, on SDS-PAGE under reducing conditions. |
| Formulation | Lyophilized after extensive dialysis against PBS. |
| Reconstitution | It is recommended that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. Reconstitute the lyophilized powder in ddH ₂ O or PBS up to 100 μg/ml. |
| Storage & Stability | Upon receiving, this product remains stable for up to 6 months at lower than -70°C. Upon reconstitution, the product should be stable for up to 1 week at 4°C or up to 3 months at -20°C. For long term storage it is recommended that a carrier protein (example 0.1% BSA) be added. Avoid repeated freeze-thaw cycles. |

Background

Target Background : Chemokine (C-C motif) ligand 1 (CCL1), also known as I-309, is a small glycoprotein secreted by activated T cells that belongs to the family of chemokines. Human CCL1 has been assumed to be a homologue of mouse TCA3. While the two proteins share only approximately 42% amino acid sequence identity, both chemokines contain an extra pair of cysteine residues not found in most other chemokines. CCL1 attracts monocytes, NK cells, immature B cells and dendritic cells by interacting with the cell surface chemokine receptor CCR8. This chemokine resides in a large cluster of CC chemokines on human chromosome 17.



Synonyms: CCL-1; I-309; TCA-3

For laboratory research use only. Direct human use, including taking orally and injection and clinical use are forbidden.