

Rev04
Update: Mar,01,2022

DATASHEET

FGF-basic (145aa), Human

Cat. No.: Z03310

Product Introduction

Species	Human
Protein Construction	FGF-basic (Ala144-Ser288) Accession # P09038
Purity	> 95% as analyzed by SDS-PAGE
Endotoxin Level	< 0.2 EU/μg of protein by gel clotting method
Biological Activity	ED ₅₀ < 0.25 ng/ml, measured by the cell proliferation assay using 3T3 cells, corresponding to a specific activity of > 4.0 × 10 ⁶ units/mg
Expression System	E. coli
Apparent Molecular Weight	~16.4 kDa, on SDS-PAGE under reducing conditions.
Formulation	Lyophilized after extensive dialysis against PBS.
Reconstitution	It is recommended that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. Reconstitute the lyophilized powder in ddH ₂ O or PBS up to 100 μg/ml.
Storage & Stability	Upon receiving, this product remains stable for up to 6 months at lower than -70°C. Upon reconstitution, the product should be stable for up to 1 week at 4°C or up to 3 months at -20°C. For long term storage it is recommended that a carrier protein (example 0.1% BSA) be added. Avoid repeated freeze-thaw cycles.

Background

Target Background : Fibroblast Growth Factor-basic (FGF-basic), also known as FGF-2, is a pleiotropic cytokine and one of the prototypic members of the heparin-binding FGF family. Like other FGF family members, FGF-basic has the β trefoil structure. In vivo, FGF-basic is produced by a variety of cells, including cardiomyocytes, fibroblasts, and vascular cells. FGF-basic regulates a variety of processes including cell proliferation, differentiation, survival, adhesion, motility, apoptosis, limb formation and wound healing. FGF-basic can be tumorigenic due to its role in angiogenesis and blood vessel remodeling. The angiogenic effects of FGF-basic can produce beneficial cardioprotection during acute heart injury.

Synonyms : Fibroblast Growth Factor-basic; FGF-2; BFGF; FGFB; HBGF-2; bFGF; Prostatropin

For laboratory research use only. Direct human use, including taking orally and injection and clinical use are forbidden.