

Rev03
Update: Dec,14,2021

DATASHEET

IL-17 RD, Mouse

Cat. No.: Z03328

Product Introduction

Species	Mouse
Protein Construction	IL-17 RD (Gly28-Arg299) Accession # Q8JZL1-1
Purity	> 95% as analyzed by SDS-PAGE > 95% as analyzed by HPLC
Endotoxin Level	< 0.2 EU/μg of protein by gel clotting method
Biological Activity	ED ₅₀ < 5.0 μg/ml, measured in a cell proliferation assay using wt MEF cells.
Expression System	CHO
Apparent Molecular Weight	53~56 kDa, on SDS-PAGE under reducing conditions.
Formulation	Lyophilized from a 0.2 μm filtered solution in PBS.
Reconstitution	It is recommended that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. Reconstitute the lyophilized powder in ddH ₂ O or PBS up to 100 μg/ml.
Storage & Stability	Upon receiving, this product remains stable for up to 6 months at lower than -70°C. Upon reconstitution, the product should be stable for up to 1 week at 4°C or up to 3 months at -20°C. For long term storage it is recommended that a carrier protein (example 0.1% BSA) be added. Avoid repeated freeze-thaw cycles.

Background

Target Background : Interleukin-17 receptor D (IL-17RD) , also known as similar expression to the fibroblast growth factor receptor (SEF), is a member of the interleukin-17 receptor family. Studies indicate that IL-17RD interacts with IL-17R to facilitate IL-17 signaling. Additionally, IL-17RD may play an important role in preventing certain types of cancer. Studies suggest that IL-17D acts as a feedback inhibitor of fibroblast growth factor-mediated MAPK signaling and ERK activation. Studies indicate that IL-17D can bind to MEK/ERK complexes preventing nuclear ERK translocation and subsequent cell proliferation.

Synonyms : IL17RLM; SEF

For laboratory research use only. Direct human use, including taking orally and injection and clinical use are forbidden.