

Rev04
Update: Jan,19,2024

DATASHEET

TNF- α (80-235aa), Mouse


Cat. No.: Z03333

Product Introduction


Species	Mouse
Protein Construction	TNF-α (Leu80-Leu235) Accession # P06804
Purity	> 98% as analyzed by SDS-PAGE > 98% as analyzed by HPLC
Endotoxin Level	< 0.2 EU/ μ g of protein by gel clotting method
Biological Activity	ED ₅₀ < 0.05 ng/ml, measured in a cytotoxicity assay using L-929 mouse fibrosarcoma cells, corresponding to a specific activity of > 2.0 \times 10 ⁷ units/mg.
Expression System	E. coli
Apparent Molecular Weight	~17.3 kDa, on SDS-PAGE under reducing conditions.
Formulation	Lyophilized after extensive dialysis against PBS.
Reconstitution	It is recommended that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. Reconstitute the lyophilized powder in ddH ₂ O or PBS up to 100 μ g/ml.
Storage & Stability	Upon receiving, this product remains stable for up to 6 months at lower than -70°C. Upon reconstitution, the product should be stable for up to 1 week at 4°C or up to 3 months at -20°C. For long term storage it is recommended that a carrier protein (example 0.1% BSA) be added. Avoid repeated freeze-thaw cycles.

Examples

ED₅₀ < 0.05 ng/ml, measured in a cytotoxicity assay using L-929 mouse fibrosarcoma cells, corresponding to a specific activity of > 2.0 × 10⁷ units/mg.



Lane 1: 2 μg of TNF-α (80-235aa), Mouse, non-reducing (NR)
 Lane 2: 2 μg of TNF-α (80-235aa), Mouse, reducing (R)
 > 98% as analyzed by SDS-PAGE



Background

Target Background : Tumor Necrosis Factor-Alpha (TNF-alpha) plays a major role in growth regulation, differentiation, inflammation, viral replication, tumorigenesis, and autoimmune disease. Besides inducing hemorrhagic necrosis of tumors, TNF has been found to be involved in tumorigenesis, tumor metastasis, viral replication, septic shock, fever, inflammation, and autoimmune diseases such as Crohn's disease, rheumatoid arthritis and graft-versus-host disease. TNF alpha-1a is a potent lymphoid factor that exerts cytotoxic effects on a wide range of tumor cells and certain other target cells.

Synonyms : TNF-alpha; Tumor necrosis factor ligand superfamily member 2; TNF-a; Cachectin; DIF; TNFA; TNFSF2

For laboratory research use only. Direct human use, including taking orally and injection and clinical use are forbidden.

生产商: 南京金斯瑞生物科技有限公司 江苏省南京市江宁区科学园雍熙路28号

Manufacturer: Nanjing GenScript Biotech Co., Ltd. No. 28Yongxi Road, Jiangning District, Nanjing, Jiangsu, China