

Rev04
Update: Mar,01,2022

DATASHEET


B7-H2/ICOSLG His, Human

Cat. No.: Z03415

Product Introduction

Species	Human				
Protein Construction	<table><tr><td>B7-H2/ICOSLG (Asp19-Ser258) Accession # 075144</td><td>Poly-His</td></tr><tr><td>N-term</td><td>C-term</td></tr></table>	B7-H2/ICOSLG (Asp19-Ser258) Accession # 075144	Poly-His	N-term	C-term
B7-H2/ICOSLG (Asp19-Ser258) Accession # 075144	Poly-His				
N-term	C-term				
Purity	> 95% as analyzed by SDS-PAGE				
Endotoxin Level	< 0.2 EU/μg of protein by gel clotting method				
Biological Activity	Immobilized B7-H2/ICOSLG His, Human at 1.0 μg/ml (100 μl/well) can bind ICOS Fc Chimera, Human(Cat. No.: Z03412).				
Expression System	HEK 293				
Apparent Molecular Weight	45~55 kDa, on SDS-PAGE under reducing conditions.				
Formulation	Lyophilized from a 0.2 μm filtered solution in PBS, 5% trehalose and mannitol.				
Reconstitution	It is recommended that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. Reconstitute the lyophilized powder in ddH ₂ O or PBS up to 100 μg/ml.				
Storage & Stability	Upon receiving, this product remains stable for up to 6 months at lower than -70°C. Upon reconstitution, the product should be stable for up to 1 week at 4°C or up to 3 months at -20°C. For long term storage it is recommended that a carrier protein (example 0.1% BSA) be added. Avoid repeated freeze-thaw cycles.				

Examples



Immobilized B7-H2-6His, Human at 1 µg/mL (100 µL/well) can bind ICOS Fc Chimera, Human (Z03412) with a linear range of 2.29-185.2 ng/mL.

Background

Target Background : B7-H2, best known as the ligand of inducible costimulator, belongs to B7-CD28 family, is a transmembrane glycoprotein of approximately 60 kDa. B7-H2 is expressed on antigen presenting cells such as B cells, macrophages, dendritic cells, and also in monocytes, and is a ligand for CD28 and CTLA-4 in human, whereas these interactions are not conserved in mouse. B7-H2 and B7-1 or B7-2 interacted with CD28 through distinctive domains. B7-H2-CD28 interaction is essential for the costimulation of human T cells' primary responses to allogeneic antigens and memory recall responses.

Synonyms : ICOSLG; B7-H2; B7H2; B7RP-1; B7RP1; CD275; GL50; ICOS-L; ICOSL; LICOS; ICOS ligand

For laboratory research use only. Direct human use, including taking orally and injection and clinical use are forbidden.