

Rev04
 Update: Mar,01,2022

DATASHEET

FGF-12, Human

Cat. No.: Z03573

Product Introduction

Species	Human
Protein Construction	<div style="display: flex; align-items: center; gap: 10px;"> <div style="background-color: #0056b3; color: white; padding: 5px; text-align: center;"> FGF-12 (Met1-Thr181) Accession # P61328-2 </div> <div style="background-color: #76b82a; color: white; padding: 5px; text-align: center;"> Poly-His </div> </div> <div style="display: flex; justify-content: space-around; margin-top: 5px; font-size: small;"> N-term C-term </div>
Purity	> 95% as analyzed by SDS-PAGE
Endotoxin Level	< 1 EU/μg of protein by LAL method
Biological Activity	Immobilized Human FGF-12 at 2.0 ug/ml can bind Human FGFR3-Fc. The ED ₅₀ of Human FGF-12 is 0.5-4.0 μg/ml.
Expression System	E. coli
Apparent Molecular Weight	~19 kDa, on SDS-PAGE under reducing conditions.
Formulation	Lyophilized from a 0.2 μm filtered solution in 20 mM PB, 150 mM NaCl, 5 mM EDTA, pH 7.5.
Reconstitution	It is recommended that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. Reconstitute the lyophilized powder in distilled water up to 100 μg/ml.
Storage & Stability	Upon receiving, this product remains stable for up to 6 months at -70°C or -20°C. Upon reconstitution, the product should be stable for up to 1 week at 4-7°C and up to 3 months at -20 °C or below. Avoid repeated freeze-thaw cycles.

Background

Target Background : Fibroblast Growth Factor 12 (FGF-12) is a member of the fibroblast growth factor (FGF) family. FGF-12 is probably involved in nervous system development and function. FGF-12 lacks the N-terminal signal sequence present in most of the FGF family members, but it contains clusters of basic residues that have been demonstrated to act as a nuclear localization signal. When transfected into mammalian cells, this protein accumulated in the nucleus, but was not secreted. The specific function of this gene has not yet been determined. Two alternatively spliced transcript variants encoding distinct isoforms have been reported.

Synonyms : Fibroblast Growth Factor 12; Fibroblast Growth Factor Homologous Factor 1; FHF-1; Myocyte-Activating Factor; FGF12; FGF12B; FHF1

For laboratory research use only. Direct human use, including taking orally and injection and clinical use are forbidden.